

AMENDMENTS TO THE CLAIMS:

Please cancel claim 27, without prejudice.

Please amend claims 22 and 24 to correct typographical errors.

1. (Withdrawn) A conditional replication shuttle vector comprising:
 - (a) an R6K origin of replication; and
 - (b) a nucleic acid encoding a recombination protein.
2. (Withdrawn) The conditional replication shuttle vector of Claim 1 wherein the recombination protein is recA
3. (Withdrawn) The conditional replication shuttle vector of Claim 1 further comprising a nucleic acid encoding a marker protein.
4. (Withdrawn) The conditional replication shuttle vector of Claim 3 wherein the nucleic acid encoding the marker protein is IRES-EGFP.
5. (Withdrawn) The conditional replication shuttle vector of Claim 3 further comprising a second marker protein.
6. (Withdrawn) The conditional replication shuttle vector of Claim 5 wherein the nucleic acid encoding the second marker protein is taulacZ.
7. (Withdrawn) The conditional replication shuttle vector of Claim1 further comprising a gene that can be counter-selected against.
8. (Withdrawn) The conditional replication shuttle vector of Claim 7 wherein the gene that can be counter-selected against is SacB.
9. (Withdrawn) The conditional replication shuttle vector of Claim 7 wherein the gene that can be counter-selected against confers tetracycline resistance.

10. (Withdrawn) The conditional replication shuttle vector of Claim 1 further comprising an A box region bracketed by two restriction enzyme sites; wherein said A box region and said restriction enzyme sites can be used to insert a selected nucleic acid into said conditional replication shuttle vector.

11. (Withdrawn) The conditional replication shuttle vector of Claim10 wherein the two restriction enzyme sites are AscI and SmaI.

12. (Withdrawn) The conditional replication shuttle vector of Claim1 further comprising two FRT sites; wherein the two FRT sites are on opposite sides of the A box.

13. (Withdrawn) The conditional replication shuttle vector of Claim 1 further comprising two homologous nucleotide sequences of 500 basepairs or more; wherein the two homologous nucleotide sequences are on opposite sides of the A box.

14. (Withdrawn) The conditional replication shuttle vector of Claim13 wherein the two homologous nucleotide sequences encode the enhanced green fluorescent protein (EGFP).

15. (Previously Presented) A method of selectively performing homologous recombination with a particular nucleotide sequence of a Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC) that is contained in a recombination deficient host cell comprising introducing a conditional replication shuttle vector into a recombination deficient host cell and therein enabling homologous recombination in the host cell via the transient expression of a recombination protein in the host cell;

wherein the host cell comprises a Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC) which contains the particular nucleotide sequence; wherein the conditional replication shuttle vector encodes a recombination protein that is transiently expressed by the host cell; wherein the conditional replication shuttle vector contains homologous nucleic acid sequences capable of selectively integrating into the particular nucleotide sequence when the recombination protein is expressed; and wherein the expressed recombination protein effectuates recombination of the shuttle vector and the Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC).

16. (Previously Presented) A method of selectively modifying a particular nucleotide sequence of a Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC) that is contained in a recombination deficient host cell comprising:

(a) introducing a conditional replication shuttle vector into a recombination deficient host cell; wherein the host cell comprises a Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC) that comprises a gene of interest which contains the particular nucleotide sequence; wherein the conditional replication shuttle vector encodes a recombination protein that is expressed by the host cell and permits homologous recombination to occur in the host cell; wherein the conditional replication shuttle vector contains homologous nucleic acid sequences capable of selectively integrating into the particular nucleotide sequence when the recombination protein is expressed forming a co-integrate; wherein the nucleic acid sequences that selectively integrate into the particular nucleotide sequence and the nucleic acid encoding the recombination protein are positioned on the conditional replication shuttle vector such that upon resolution of the co-integrate, the nucleic acid encoding the recombination protein remains with the conditional replication shuttle vector; and wherein the expressed recombination protein effectuates recombination of the shuttle vector and the Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC); and

(b) growing the host cell under conditions in which the conditional replication shuttle vector cannot replicate, therein diluting out the conditional replication shuttle vector encoding the recombination protein, and thereby preventing further recombination events in the recombination deficient cells.

17. (Original) The method of Claim 16 wherein the conditional replication shuttle vector further comprises a nucleic acid that encodes a marker protein or peptide and wherein the nucleic acid that selectively integrates into the particular nucleotide sequence and the nucleic acid encoding the marker protein or peptide are positioned on the conditional replication shuttle vector such that upon resolution of the co-integrate, the nucleic acid encoding the marker protein or peptide is inserted into or adjacent to the particular nucleotide sequence.

18. (Previously Presented) The method of Claim 16 wherein the conditional replication shuttle vector cannot replicate in the host cell because the conditional replication shuttle vector requires a particular protein for replication and neither the host cell nor the Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC) encode the particular protein.

19. (Canceled)

20. (Previously Presented) The method of Claim 16 wherein the BBPAC is a BAC or a PAC.

21. (Original) The method of Claim 20 wherein the conditional replication shuttle vector cannot replicate in the host cell because the conditional replication shuttle vector comprises a R6K origin of replication and neither the host cell nor the BAC encode *pir*.

22. (Currently Amended) The method of Claim 21 wherein the conditional replication shuttle vector further comprises a first *frt* site that is positioned on one side of the nucleic acid that selectively integrates into the particular nucleotide sequence, and a second *frt* site that is positioned on the other side of the nucleic acid that selectively integrates into the particular nucleotide sequence and wherein the resolution of the co-integrate is performed by adding ~~flp~~ flp recombinase to the host cell.

23. (Original) The method of Claim 21 wherein the conditional replication shuttle vector further comprises a nucleic acid encoding a marker protein or peptide that is positioned in between the two *frt* sites and is also adjacent to the nucleic acid that selectively integrates into the particular nucleotide sequence such that after the resolution, the marker protein or peptide is contained by the BAC.

24. (Currently Amended) The method of Claim 22 wherein ~~flp~~ flp recombinase is added to the host cell by introducing a plasmid that encodes ~~flp~~ flp recombinase to the host cell.

25. (Original) The method of Claim 24 wherein the plasmid contains a conditional origin of replication.

26. (Original) The method of Claim 21 wherein the conditional replication shuttle vector further comprises two homologous nucleotide sequences that are homologous to each other but are not homologous to the BAC; wherein the two homologous nucleotide sequences are positioned on the conditional replication shuttle vector to be on opposite sides of the nucleic acid that selectively integrates into the particular nucleotide sequence; and wherein the resolution of the co-integrate is performed by a recombination event between the two homologous nucleotide sequences.

27. (Canceled)

28. (Original) The method of Claim 21 wherein the recombination deficient host cell cannot independently support homologous recombination because the host cell is *RecA⁻*.

29. (Original) The method of Claim 21 further comprising adding a counterselection agent after the resolution of the co-integrate to remove host cells that comprise the conditional replication shuttle vector; wherein the conditional replication shuttle vector further comprises a counterselection gene that is positioned on the conditional replication shuttle vector such that upon resolution of the co-integrate the counterselection gene remains with the conditional replication shuttle vector.

30. (Original) The method of Claim 29 wherein the counterselection agent is sucrose and the counterselection gene is SacB.

31. (Original) The method of Claim 30 wherein the recombination deficient host cell cannot independently support homologous recombination because the host cell is RecA⁻.

32. (Original) The method of Claim 31 wherein the recombination protein is selected from the group consisting of recA, the rec E and rec T protein pair, the Lambda *beta* protein, and the *Arabidopsis thaliana* DRT100 gene product.

33. (Withdrawn) An IOBCV that comprises a particular nucleotide sequence that has been modified by the method of Claim 16.

34. (Withdrawn) A BAC that comprises a particular nucleotide sequence that has been modified by the method of Claim 21.

35. (Withdrawn) A BAC that comprises a particular nucleotide sequence that has been modified by the method of Claim 26.

36. (Withdrawn) A method of producing a non-human transgenic animal comprising:

(a) introducing a Bacterial or Bacteriophage-Derived Artificial Chromosome (BBPAC) into a eukaryotic cell; and

(b) placing the eukaryotic cell into a recipient animal, wherein the eukaryotic cell develops into the non-human transgenic animal;

wherein the BBPAC has been modified through homologous recombination in a RecA⁻ bacterial host cell that has been induced to support homologous recombination by the transient expression of a

recombination protein; and wherein the eukaryotic cell is selected from the group consisting of a fertilized animal zygote and an embryonic stem cell.

37. (Withdrawn) The method of Claim 36 wherein said eukaryotic cell is a fertilized animal zygote and said introducing is performed by pronuclear injecting the BBPAC into the fertilized animal zygote.

38. (Withdrawn) The method of Claim 37 wherein the BBPAC is a Bacterial Artificial Chromosome (BAC); the animal is a mouse; and the fertilized animal zygote is a C57BL/6 mouse zygote.

39. (Withdrawn) The method of Claim 38 wherein said eukaryotic cell is a mouse embryonic stem (ES) cell and said introducing is performed by transfecting the mouse ES cell.

40. (Withdrawn) The method of Claim 39 wherein the BBPAC is a Bacterial Artificial Chromosome (BAC).

41. (Withdrawn) The method of Claim 40 wherein the animal is a mammal.

42. (Withdrawn) The method of Claim 41 wherein the mammal is a mouse.

43. (Withdrawn) A method of producing a non-human transgenic animal comprising:

- (a) introducing the BAC of Claim 34 into a eukaryotic cell; and
- (b) placing the eukaryotic cell into a recipient animal, wherein the eukaryotic cell develops into the non-human transgenic animal.

44. (Withdrawn) A method of producing a non-human transgenic animal comprising:

- (a) introducing the BAC of Claim 35 into a eukaryotic cell; and
- (b) placing the eukaryotic cell into a recipient animal, wherein the eukaryotic cell develops into the non-human transgenic animal.

45. (Withdrawn) A non-human transgenic animal obtained by the method of Claim 44.

46. (Withdrawn) A non-human transgenic animal obtained by the method of Claim 43.

47. (Withdrawn) A non-human transgenic animal obtained by the method of Claim 36.

48. (Previously Presented) The method of claim 16, wherein said BBPAC is modified after resolution to result in nucleic acids insertions, and/or nucleic acids deletions, and/or point mutations on the BBPAC.